Periodic Solutions for Nonlinear Evolution Equations

Y. Q. Chen¹, Y. J. Cho² and L. Yang³

¹Department of Mathematics
Fushan Institute of Science and Technology
Fushan, Guangdong 528000, P. R. China
²Department of Mathematics
Gyeongsang National University
Chinju 660-701, Korea
³Department of Mathematics
Mianyang Teacher's Colloge
Mianyang, Sichuan 621000, P. R. China

Abstract. In this paper, we study the existence of periodic solutions of following type evolution equations

$$f(t, u(t)) \in -au''(t) + bu'(t) + A(t)u(t), \quad t \in R,$$

in infinite dimensional spaces. Our main results generalize the corresponding results in Browder [6], Hirano [16], Shioji [25] and Vrabie [26].

Keywords. Periodic solution, class (S_+) , evolution equation, topological dgree.

AMS (MOS) subject classification: 1991 Mathematics Subject Classification: Primary 34C25, 34G20; Secondary 47H05, 47H10.

1 Introduction

The purpose of this paper is to study the existence of periodic solutions of the following type evolution equation

$$f(t, u(t)) \in -au''(t) + bu'(t) + A(t)u(t), \quad t \in R,$$
 (E1.1)

in real reflexive Banach spaces or Hilbert spaces. First, in Section 2, we set a=0,b=1 and study the periodic solution of the first order evolution. Periodic solutions of the first order equations have been extensively studied by many authors. For the operator A in linear cases, see, Amann [1], Becker [5], Pruss [23], etc., and, for the operator A in nonlinear cases, see Browder [6], Chen [11], Deimling [12], Hirano [16], Shioji [25], Vrabie [26], etc.

In this paper, we consider that A is a multi-valued operator of the class (S_+) which was introduced by Petryshyn [22]. This condition is different from

²The corresponding author: Yeol Je Cho.

¹The first author is partially supported by a research fund for return from oversea of Education Ministry, P. R. China, and the second author is supported by the KOSEF 2000-1-10100-003-3.